

THE DERIVED SERIES OF A FINITE p -GROUP

CSABA SCHNEIDER

ABSTRACT. Let G be a finite p -group, and let $G^{(d)}$ denote the d -th term of the derived series of G . We show, for $p \geq 5$, that $G^{(d)} \neq 1$ implies $\log_p |G| \geq 2^d + 3d - 6$, and hence we improve a recent result by Mann.

1. INTRODUCTION

In a group G , let $G^{(d)}$ denote the d -th term of the derived series (as usual, we write $G^{(0)} = G$, $G^{(1)} = G'$, $G^{(2)} = G''$). The main goal of this article is to give a new lower bound for the order of a p -group in which $G^{(d)} \neq 1$.

Theorem 1.1. *Let p be a prime such that $p \geq 5$, and let G be a finite p -group. If $G^{(d)} \neq 1$ then $\log_p |G| \geq 2^d + 3d - 6$.*

The problem of bounding the order of a finite p -group in terms of its derived length has a long history starting from Burnside's papers [Bur13, Bur14]. It is well-known that non-abelian p -groups have order at least p^3 and non-metabelian p -groups have order at least p^6 (for $p = 2, 3$, the sharp bound is p^7). In his 1934 paper [Hal34], P. Hall noticed, for $i \geq 0$, that $G^{(i+1)} \neq 1$ implies $\log_p |G^{(i)}/G^{(i+1)}| \geq 2^i + 1$. He used this lower bound to prove that

$$(1) \quad G^{(d)} \neq 1 \quad \text{implies} \quad \log_p |G| \geq 2^d + d.$$

Recently, there has been some interest in obtaining sharper versions of Hall's theorem. Mann [Man00] showed that if G is a p -group then $G^{(d)} \neq 1$ implies $\log_p |G| \geq 2^d + 2d - 2$, and Theorem 1.1 in this paper improves the linear term of Mann's bound. Another direction in this research is to consider the assertion (1) for a fixed d . Hall's bound is,

Date: February 2, 2008.

2000 Mathematics Subject Classification. 20D15, 20-04.

Key words and phrases. finite p -groups, derived subgroups, derived quotients, derived series, associated Lie ring.

in general, sharp for $d = 1, 2$ (we mentioned above that, for $p = 2, 3$ and $d = 2$, the sharp bound is p^7), but it is not sharp already for $d = 3$. In this case, Hall's lower bound is p^{11} , but, for $p \geq 3$, the smallest p -groups in which $G^{(3)} \neq 1$ have order p^{14} (see [ER00, ERNS99, ERNXX]) and the smallest such 2-groups have order 2^{13} [ERNXX]. For $p \geq 5$ and $d \geq 3$, the smallest known groups with $G^{(d)} \neq 1$ have composition length $2^{d+1} - 2$ (see [ERNS99]), and so there is a gap to be filled.

Similar questions arise in the context of soluble groups. Glasby [GlaXX, GlaYY] studied soluble groups with a fixed derived length and smallest possible composition length.

We prove Theorem 1.1 by studying the abelian and metabelian quotients of the derived series of a finite p -group and obtain several results that are interesting in their own right. By Hall's lemma referred to above, we have, in a p -group G , that $G^{(d+1)} \neq 1$ implies $\log_p |G^{(d)}/G^{(d+1)}| \geq 2^d + 1$. In a finite p -group G we call $G^{(d)}/G^{(d+1)}$ a *small derived quotient* if

$$(2) \quad G^{(d+1)} \neq 1 \quad \text{and} \quad \log_p |G^{(d)}/G^{(d+1)}| = 2^d + 1.$$

We do not know whether, for an arbitrary p and d , the quotient $G^{(d)}/G^{(d+1)}$ can be small in a p -group G . Groups in which G/G' or G'/G'' is small are easy to construct; odd-order groups with small second derived quotients G'/G'' were characterised in [Sch03]. The Sylow 2-subgroup P of the symmetric group S_{2^d} of rank 2^d satisfies $\log_2 |P^{(d-2)}/P^{(d-1)}| = 2^{d-2} + 1$ and $P^{(d-1)} \neq 1$ (see [KLGP97, Lemma (II.7)]). Therefore small derived quotients do indeed exist. On the other hand, for $p \geq 3$ and $d \geq 2$, we do not have examples in which $G^{(d)}/G^{(d+1)}$ is small. We do, however, know that the structure of a finite p -group in which $G^{(d)}/G^{(d+1)}$ is small must be severely restricted. Blackburn in [Bla87] attributes a theorem to P. Hall that, for odd primes, if G'/G'' is small, then $|G''| = p$; see [Bla87]. In the same article, Blackburn himself shows that the same condition for 2-groups implies that G'' is a two-generator abelian group and is nearly homocyclic. Using Bokut's ideas [Bok71], this result was generalised in my PhD Thesis [Sch00], where I showed that if $p \neq 3$ and (2) holds with $d \geq 1$, then $G^{(d)}$ has nilpotency class at most 3. Further, Mann [Man00] showed that a finite p -group can have at most two small derived quotients.

It is explained in Section 3 that if (2) holds in a finite p -group G then $G^{(d)}/[G^{(d)}, G]$ must be elementary abelian with rank at most 2. In this article, I prove the following theorem, which claims that p^2 is usually not possible.

Theorem 1.2. *Suppose that p is an odd prime, $d \geq 1$, and G is a finite p -group such that $G^{(d+1)} \neq 1$. If $\log_p |G^{(d)}/G^{(d+1)}| = 2^d + 1$, then $|G^{(d)}/[G^{(d)}, G]| = p$.*

Theorem 1.2 is a generalisation of [Sch03, Theorem 1.1]. Its proof will be presented in Section 5 using the Lie algebra constructed in Section 4. In a separate article [SchXX], I use Theorem 1.2 to show, for odd p , that two distinct small derived quotients can occur only in a group with order p^6 . This gives an improvement of Mann's related result in [Man00].

Theorem 1.2 is used to obtain the constant term “6” in Theorem 1.1. In order to obtain the coefficient “3” of the linear term, we study metabelian quotients of the derived series, that is, we consider the factor groups $G^{(i)}/G^{(i+2)}$. This way we are able to prove a better lower bound for the order of G than those obtained by considering only the abelian quotients $G^{(i)}/G^{(i+1)}$.

The structure of this paper is as follows. In Section 2 we summarise the technical tools that we need in the proofs of the main theorems. We prove Theorem 1.2 in Sections 3–5. Many of the proofs in this paper use the Lie ring method. In order to verify Theorem 1.2 we define a filtration on $G^{(d)}$ and consider the graded Lie ring associated with this filtration. We identify a small subring in this Lie ring, and relatively simple, though rather lengthy, computation yields Theorem 1.2. Then, in Section 6, we study the metabelian quotients of the derived series of a finite p -group, and prove the somewhat technical Theorem 6.1. Again, the hard part is to identify a small subring in the Lie ring associated with the lower central series of G . Once this subring is found, elementary, but somewhat cumbersome, calculations leads to Theorem 6.1. Finally, in Section 7 we present a proof for Theorem 1.1.

2. A TOOLBOX

In this section we review a couple of well-known results that are used in this paper. The construction of the Lie ring in Section 4 is based on some simple properties of group commutators. In addition to the best-known commutator identities (see for instance [LGM02, Proposition 1.1.6]), the following collection formula will often be used: if x and y are elements of a finite p -group then

$$(3) \quad [x^p, y] \equiv [x, y]^p \pmod{H^p \gamma_p(H)} \quad \text{where} \quad H = \langle x, [x, y] \rangle;$$

see [LGM02, Proposition 1.1.32]. The Hall-Witt identity will occur in a lesser-known form: if x , y , and z are elements of a group then

$$(4) \quad [x, y, z^x][z, x, y^z][y, z, x^y] = 1;$$

see [LGM02, Proposition 1.1.6(v)].

The following result was proved by P. Hall [Hal34].

Lemma 2.1. *Suppose that H is a non-abelian normal subgroup in a finite p -group G such that $H \leq \gamma_i(G)$. Then $|H/H'| \geq p^{i+1}$ and $|H| \geq p^{i+2}$.*

For the proof of the following lemma, see [Bla58, Lemma 2.1].

Lemma 2.2. *If G is a group and H is a normal subgroup with cyclic quotient, then $G' = [G, H]$.*

If U and V are subgroups in a group and n is a natural number then let $[U, {}^nV]$ denote the left-normed commutator subgroup

$$[U, {}^nV] = [U, \underbrace{V, \dots, V}_{n \text{ copies}}].$$

The following result is easy to verify by induction on i .

Lemma 2.3. *If A , B are normal subgroups of a group G , then $[A, \gamma_i(B)] \leq [A, {}^iB]$.*

3. A FILTRATION ON $G^{(d)}$

In this section we start the preparation for the proof of Theorem 1.2. Let G be a finite p -group. Let us define a filtration on $G^{(d)}$ as follows:

$$N_{2^d} = G^{(d)} \quad \text{and} \quad N_{k+1} = [N_k, G] \quad \text{for} \quad k \geq 2^d.$$

Using the notation introduced in the previous section, we may write $N_k = [G^{(d)}, {}^{k-2^d}G]$. The following lemma shows, in the terminology of [Sha95], that the N_i form a strongly central filtration.

Lemma 3.1. *The filtration $\{N_k\}_{k \geq 2^d}$ satisfies the property $[N_j, N_k] \leq N_{j+k}$ for each j , $k \geq 2^d$.*

Proof. As N_k is contained in $\gamma_k(G)$, and, for a subgroup H , the commutator subgroup $[H, \gamma_i(G)]$ is contained in $[H, {}^iG]$ (see Lemma 2.3), we obtain

$$[N_j, N_k] \leq [N_j, \gamma_k(G)] \leq [N_j, {}^kG] = N_{j+k}.$$

□

Lemma 3.2. *Suppose that the derived quotient $G^{(d)}/G^{(d+1)}$ is small in a finite p -group G . Then the following is true.*

- (a) *Exactly one of the following must hold:*
 - (i) $|N_i/N_{i+1}| = p$ for $i \in \{2^d, \dots, 2^{d+1}\}$, and $G^{(d+1)} = N_{2^{d+1}+1}$;
 - (ii) $|G^{(d)}/N_{2^{d+1}}| = p^2$, $|N_i/N_{i+1}| = p$ for $i \in \{2^d + 1, \dots, 2^{d+1} - 1\}$, and $G^{(d+1)} = N_{2^{d+1}}$.
- (b) N_i/N_{i+1} is elementary abelian for $i \geq 2^d$.
- (c) $N_{2^{d+1}} = G^{(d)} \cap \gamma_{2^{d+1}}(G)$.

Proof. (a) As $G^{(d)} \leq \gamma_{2^d}(G)$, we obtain

$$G^{(d+1)} = [G^{(d)}, G^{(d)}] \leq [G^{(d)}, \gamma_{2^d}(G)] \leq [G^{(d)}, {}^{2^d}G] = N_{2^{d+1}},$$

therefore we have the following chain of G -normal subgroups:

$$(5) \quad G^{(d)} > [G^{(d)}, G] > [G^{(d)}, G, G] > \dots > [G^{(d)}, {}^{2^d}G] \geq G^{(d+1)}.$$

Counting number of non-trivial factors of this chain, we obtain that $G^{(d)}/[G^{(d)}, G]$ has order at most p^2 . If $G^{(d)}/[G^{(d)}, G]$ is cyclic, then, by Lemma 2.2, the subgroup $G^{(d+1)}$ coincides with $[G^{(d)}, [G^{(d)}, G]]$, and so

$$G^{(d+1)} = [G^{(d)}, G^{(d)}] = [G^{(d)}, [G^{(d)}, G]] \leq [G^{(d)}, \gamma_{2^{d+1}}(G)] \leq [G^{(d)}, {}^{2^{d+1}}G] = N_{2^{d+1}+1}.$$

Thus, in this case, we obtain the following modified chain:

$$(6) \quad G^{(d)} > [G^{(d)}, G] > [G^{(d)}, G, G] > \dots > [G^{(d)}, {}^{2^d}G] > [G^{(d)}, {}^{2^{d+1}}G] \geq G^{(d+1)}.$$

If the first quotient in these chains has order p , then this quotient is cyclic, and so (6) must hold. In this case, counting the non-trivial factors in (6), we find that case (i) must be valid. Now suppose that the first quotient $G^{(d)}/[G^{(d)}, G]$ has order p^2 . Then chain (6) is too long, and so $G^{(d)}/[G^{(d)}, G]$ must be elementary abelian. As before, we count the number of factors in (5) and find that case (ii) must hold.

(b) In the previous paragraph we obtained that $G^{(d)}/[G^{(d)}, G]$ is elementary abelian, and we use induction to show that so are the other factors N_i/N_{i+1} . Let us suppose that

N_{k-1}/N_k is elementary abelian for some $k \geq 2^d + 1$. Consider a generator $a = [x, y]$ of N_k where $x \in N_{k-1}$ and $y \in G$. Then $[x, y]^p \equiv [x^p, y]$ modulo N_{k+1} (equation (3)) and x^p lies in N_k by the induction hypothesis. Hence we obtain that $[x^p, y] \in N_{k+1}$, and so $[x, y]^p \in N_{k+1}$ as required.

(c) As $G^{(d)} \leq \gamma_{2^d}(G)$, and $N_{2^d+1} = [G^{(d)}, G]$, we obtain $N_{2^d+1} \leq G^{(d)} \cap \gamma_{2^d+1}(G)$. So it suffices to prove that $G^{(d)} \cap \gamma_{2^d+1}(G) \leq N_{2^d+1}$. Assume now that this claim is false, and choose $x \in G^{(d)} \cap \gamma_{2^d+1}(G)$ such that $x \notin N_{2^d+1}$. Let M be the subgroup $\langle N_{2^d+1}, x \rangle$. Note that M is contained in $\gamma_{2^d+1}(G)$ and that N_{2^d+1} is a proper subgroup of M . If case (i) of part (a) is valid, then $M = G^{(d)}$. Thus, in this case, $G^{(d)} \leq \gamma_{2^d+1}(G)$, and so Lemma 2.1 implies that $G^{(d)}/G^{(d+1)}$ cannot be a small derived quotient. If case (ii) of part (a) is valid, then, as $G^{(d)}/N_{2^d+1}$ is elementary abelian with order p^2 , M is a maximal subgroup in $G^{(d)}$, and hence $G^{(d)}/M$ is cyclic. Therefore, by Lemma 2.2, $G^{(d+1)} = [G^{(d)}, G^{(d)}] = [G^{(d)}, M]$. Since M is contained in $\gamma_{2^d+1}(G)$, $G^{(d+1)}$ is contained in $[G^{(d)}, \gamma_{2^d+1}(G)]$, which, in turn, lies in $[G^{(d)}, {}^{2^d+1}G] = N_{2^d+1+1}$. This, however, is a contradiction, as, in this case, $G^{(d+1)} = N_{2^d+1}$. $\square$

4. A LIE ALGEBRA ASSOCIATED WITH $G^{(d)}$

We stick to the notation introduced in the previous section.

In the following two sections we assume that G is a finite p -group, $d \geq 1$, $G^{(d)}/G^{(d+1)}$ is a small derived quotient, and that $G^{(d)}/[G^{(d)}, G]$ is elementary abelian with rank 2. Our aim is to show at the end of Section 5 that $p = 2$, and hence we prove Theorem 1.2. We may also assume without loss of generality that $|G^{(d+1)}| = p$, and so $G^{(d+1)}$ lies in the centre of G . Lemma 3.2 shows that in this case $|N_{2^d+1}| = p$, and that $N_{2^d+1+1} = 1$.

For the proof of Theorem 1.2 we introduce three Lie algebras. The first, we denote it by K , will be a graded Lie algebra associated with the filtration $\{N_i\}$ of $G^{(d)}$. The second algebra $\overline{K}$ will be an extension of K , in the sense that K will be an ideal of $\overline{K}$, and finally, L will be a small subalgebra of $\overline{K}$. We will carry out most of our computations in L .

Let K be the graded Lie ring associated with the filtration $\{N_i\}$ defined in the previous section. In other words we define K as

$$K = \bigoplus_{i=2^d}^{2^{d+1}} N_i/N_{i+1}$$

and the multiplication between two homogeneous elements of K is defined as follows:

$$[aN_{i+1}, bN_{j+1}] = [a, b]N_{i+j+1} \quad \text{for } a \in N_i, b \in N_j.$$

From Lemma 3.2 it also follows that K can be viewed as a Lie algebra over $\mathbb{F}_p$. Let K_i denote the homogeneous components of K ; that is $K_i = N_i/N_{i+1}$ for $i \geq 2^d$. It is a consequence of Lemma 3.1 that $[K_i, K_j] \leq K_{i+j}$.

Let Q denote the subgroup $[G^{(d-1)}, G](G^{(d-1)})^p$.

Lemma 4.1. *We have*

- (i) $[G^{(d-1)}, Q] \leq N_{2^{d+1}}$;
- (ii) $[N_i, Q] \leq N_{i+2^{d-1}+1}$, for $i \geq 2^d$.

Proof. (i) By [LGM02, Lemma 1.1.9(ii)], it suffices to show that

$$[G^{(d-1)}, [G^{(d-1)}, G]] \leq N_{2^{d+1}} \quad \text{and} \quad [G^{(d-1)}, (G^{(d-1)})^p] \leq N_{2^{d+1}}.$$

Since $[G^{(d-1)}, [G^{(d-1)}, G]] \leq G^{(d)} \cap \gamma_{2^{d+1}}(G)$, using Lemma 3.2(c), we obtain that $[G^{(d-1)}, [G^{(d-1)}, G]] \leq N_{2^{d+1}}$. Suppose that $x, y \in G^{(d-1)}$. Then $[x, y^p] \equiv [x, y]^p \pmod{H^p \gamma_p(H)}$ where $H = \langle y, [x, y] \rangle$ (see equation (3)). As $G^{(d)}/N_{2^{d+1}}$ has exponent p , we have that $[x, y]^p \in N_{2^{d+1}}$, and, since

$$H^p \gamma_p(H) \leq G^{(d)} \cap \gamma_{2^{d+1}}(G) = N_{2^{d+1}},$$

we obtain that $[x, y^p] \in N_{2^{d+1}}$. As $[G^{(d-1)}, (G^{(d-1)})^p]$ is generated as a normal subgroup by such elements $[x, y^p]$ (see [LGM02, Lemma 1.1.9(iii)]), we have $[G^{(d-1)}, (G^{(d-1)})^p] \leq N_{2^{d+1}}$.

(ii) As above, it suffices to show that

$$[N_i, [G^{(d-1)}, G]] \leq N_{i+2^{d-1}+1} \quad \text{and} \quad [N_i, (G^{(d-1)})^p] \leq N_{i+2^{d-1}+1}.$$

Both claims of the last displayed equation can, however, be shown as the assertions in the proof of part (i). $\square$

Set $K_{2^{d-1}}$ to be $G^{(d-1)}/Q$. Clearly $K_{2^{d-1}}$ is elementary abelian, and so it can be considered as an $\mathbb{F}_p$ -vector space. Let $\overline{K} = K_{2^{d-1}} \oplus K$. Let us define a graded Lie algebra structure on $\overline{K}$. Two elements of K are multiplied according to the product rule in K . If $aQ \in K_{2^{d-1}}$ and $bN_{i+1} \in K$ then let us define

$$[aQ, bN_{i+1}] = [a, b]N_{2^{d-1}+i+1}.$$

If $aQ, bQ \in K_{2^{d-1}}$ then we define

$$[aQ, bQ] = [a, b]N_{2^d+1}.$$

We extend this product linearly and anti-symmetrically to $\overline{K}$.

Lemma 4.2. *The product on $\overline{K}$ is well-defined, and the algebra $\overline{K}$ is a Lie algebra over $\mathbb{F}_p$.*

Proof. In order to show that this lemma is true we have to verify that the product between homogeneous elements of $\overline{K}$ satisfies the following properties:

- (i) it is well-defined;
- (ii) it is linear;
- (iii) it is anti-symmetric;
- (iv) and the Jacobi identity holds.

Note that K is a Lie ring, and so these properties obviously hold for homogeneous elements of K .

(i) We claim that the product between an element of $K_{2^{d-1}}$ and a homogeneous element of K is well-defined. We are required to show that if $x, x' \in N_j$ and $y, y' \in G^{(d-1)}$ such that $xN_{j+1} = x'N_{j+1}$ and $yQ = y'Q$, then $[xN_{j+1}, yQ] = [x'N_{j+1}, y'Q]$. This amounts to saying that

$$[x, y]N_{j+2^{d-1}+1} = [x', y']N_{j+2^{d-1}+1},$$

that is $[x, y] \equiv [x', y'] \pmod{N_{j+2^{d-1}+1}}$. There are some $n \in N_{j+1}$ and $g \in Q$ such that $x' = xn$ and $y' = yg$. Now

$$[x', y'] = [xn, yg] = [x, yg][x, yg, n][n, yg] \equiv [x, yg] = [x, g][x, y][x, y, g] \equiv [x, y]$$

where congruences are taken modulo $N_{j+2^{d-1}+1}$. Note that we used Lemma 4.1(ii). Using Lemma 4.1(i), a similar argument shows that the product is well-defined between two elements of $K_{2^{d-1}}$.

(ii) Suppose that $x_1, x_2 \in G^{(d-1)}$, and that $y \in N_j$ with some $j \geq 2^d$. Then, by Lemma 2.3, $[x_1, y, x_2] \in N_{j+2^{d-1}+1}$, and so

$$\begin{aligned} [x_1Q + x_2Q, yN_{j+1}] &= [x_1x_2Q, yN_{j+1}] = [x_1x_2, y]N_{j+2^{d-1}+1} \\ &= [x_1, y][x_1, y, x_2][x_2, y]N_{j+2^{d-1}+1} = [x_1, y][x_2, y]N_{j+2^{d-1}+1} \\ &= [x_1, y]N_{j+2^{d-1}+1} + [x_2, y]N_{j+2^{d-1}+1} = [x_1Q, yN_{j+1}] + [x_2Q, yN_{j+1}]. \end{aligned}$$

Similar argument shows that the Lie bracket is linear between homogeneous elements of $K_{2^{d-1}}$; that $[-xQ, yQ] = [xQ, -yQ] = -[xQ, yQ]$, for $x, y \in G^{(d-1)}$; and that $[-xQ, yN_{i+1}] = [xQ, -yN_{i+1}] = -[xQ, yN_{i+1}]$ for $x \in G^{(d-1)}$ and $y \in N_i$.

(iii) It is an easy computation to show that the Lie bracket is anti-symmetric.

(iv) Let us finally show that the Jacobi identity holds. Using the commutator identities, we compute

$$(7) \quad [x, y, z^x] = [x, y, z[z, x]] = [x, y, [z, x]][x, y, z][x, y, z, [z, x]].$$

If $x, y, z \in G^{(d-1)}$, then (7) implies that $[x, y, z^x] \equiv [x, y, z] \pmod{N_{2^d+2^{d-1}+1}}$. If $x, y \in G^{(d-1)}$ and $z \in N_i$, then (7) implies that $[x, y, z^x] \equiv [x, y, z] \pmod{N_{2^d+i+1}}$. Finally, if $x \in G^{(d-1)}$, $y \in N_i$, and $z \in N_j$, then $[x, y, z^x] \equiv [x, y, z] \pmod{N_{2^{d-1}+i+j+1}}$. As the Hall-Witt identity

$$[x, y, z^x][z, x, y^z][y, z, x^y] = 1$$

holds (see (4)), we find that

$$[x, y, z][z, x, y][y, z, x] \in N_m.$$

where

$$m = \begin{cases} 2^d + 2^{d-1} + 1 & \text{if } x, y, z \in G^{(d-1)}; \\ i + 2^d + 1 & \text{if } x, y \in G^{(d-1)} \text{ and } z \in N_i; \\ i + j + 2^{d-1} + 1 & \text{if } x \in G^{(d-1)}, y \in N_i, z \in N_j. \end{cases}$$

As the Jacobi identity holds for homogeneous elements of K , this shows that it also holds for homogeneous elements of $\overline{K}$. $\square$

Recall that

$$\overline{K} = K_{2^{d-1}} \oplus K = K_{2^{d-1}} \oplus K_{2^d} \oplus K_{2^{d+1}} \oplus \cdots \oplus K_{2^{d+1}}.$$

It is easy to see that $\overline{K}$ is a graded Lie algebra in the sense that $[K_i, K_j] \in K_{i+j}$ for $i, j \in \{2^{d-1}, 2^d, 2^d + 1, \dots, 2^{d+1}\}$. We define our object of interest: we let L be the subalgebra

$$L = K_{2^{d-1}} \oplus K_{2^d} \oplus K_{2^d+2^{d-1}} \oplus K_{2^{d+1}}.$$

We set $L_1 = K_{2^{d-1}}$, $L_2 = K_{2^d}$, $L_3 = K_{2^d+2^{d-1}}$ and $L_4 = K_{2^{d+1}}$. The grading on K implies that L is also graded in the sense that $[L_i, L_j] \subseteq L_{i+j}$. The following lemma lists the important properties of L .

Lemma 4.3. *The following is true:*

- (i) $\dim L_2 = 2$; $\dim L_3 = \dim L_4 = 1$;
- (ii) $L'' = L_4$;
- (iii) L is generated by L_1 .

Proof. Claim (i) follows because case (ii) of part (a) in Lemma 3.2 is valid and because we assumed that $|N_{2^{d+1}}| = p$.

Let us now prove (ii) and (iii). First we claim that $L_2 = [L_1, L_1]$. We note that $G^{(d)} = [G^{(d-1)}, G^{(d-1)}]$, and so $G^{(d)}$ is generated by elements of the form $[a, b]$ with $a, b \in G^{(d-1)}$. Hence L_{2^d} is generated by elements of the form $[a, b]N_{2^{d+1}}$. On the other hand, by the definition of the product in $\overline{K}$, $[a, b]N_{2^{d+1}} = [aQ, bQ]$, which implies that

$$L_2 = K_{2^d} = [K_{2^{d-1}}, K_{2^{d-1}}] = [L_1, L_1].$$

Since $G^{(d+1)} = [G^{(d)}, {}^2G]$ and $|G^{(d+1)}| = p$, there are elements $a, b \in G^{(d)}$ such that $[a, b] \neq 1$. The definition of K and the fact that $G^{(d+1)}$ is central implies

$$[aN_{2^{d+1}}, bN_{2^{d+1}}] = [a, b]N_{2^{d+1}+1} = [a, b],$$

which, in turn, implies that $[aN_{2^{d+1}}, bN_{2^{d+1}}] \neq 0$. As $L_2 = [L_1, L_1]$, we obtain that $aN_{2^{d+1}}, bN_{2^{d+1}} \in L'$; thus $L'' \neq 0$. As L has nilpotency class 4, and $\dim L_4 = 1$ we have that $L_4 = L''$. This completes the proof of (ii).

We have shown so far part (ii) and that $L_1, L_2, L_4 \leq \langle L_1 \rangle$. It only remains to prove that $L_3 \leq \langle L_1 \rangle$. This, however, easily follows, because

$$L_4 = L'' = [L_2, L_2] = [[L_1, L_1], [L_1, L_1]] \leq [L_1, L_1, L_1, L_1],$$

and so there must be $x_1, x_2, x_3, x_4 \in L_1$ such that $[x_1, x_2, x_3, x_4] \neq 0$. Thus $[x_1, x_2, x_3] \neq 0$. Since $[x_1, x_2, x_3] \in L_3$ and $\dim L_3 = 1$, we have $L_3 \leq \langle L_1 \rangle$. $\square$

5. THE PROOF OF THEOREM 1.2

We verify, in this section, that the assumptions made at the beginning of Section 4 imply that $p = 2$ and hence we complete the proof of Theorem 1.2.

We claim that elements $a, b, c \in L_1$ can be chosen such that the following hold:

- (i) $L_2 = \langle [a, b], [a, c] \rangle$ and $L_4 = \langle [[a, b], [a, c]] \rangle$;
- (ii) $[b, c] = [a, b, a] = [a, b, b] = [a, c, a] = 0$.

By Lemma 4.3(i), $\dim L_2 = 2$, and, by Lemma 4.3(iii), $L_2 = [L_1, L_1]$, and so there are $x, y, u, v \in L_1$ such that $L_2 = \langle [x, y], [u, v] \rangle$. If $[x, u] = [x, v] = [y, u] = [y, v] = 0$ then set $a = y + u, b = x, c = v$. Otherwise assume without loss of generality that $[x, u] \neq 0$, and so there are scalars $\alpha, \beta \in \mathbb{F}_p$ such that $[x, u] = \alpha[x, y] + \beta[u, v]$. If $\alpha \neq 0$, then we set $a = u, b = x, c = v$. If $\alpha = 0$ then $\beta \neq 0$ and we set $a = x, b = u, c = y$. It is easy to see that with each of the choices of a, b, c we have that $L_2 = \langle [a, b], [a, c] \rangle$ and that, since $L'' = L_4$, it also follows that $L_4 = \langle [[a, b], [a, c]] \rangle$.

We claim that a, b, c above can be chosen so that $[b, c] = 0$. Indeed, since $\dim L_2 = 2$, some non-trivial linear combination of $[a, b], [a, c], [b, c]$ is zero:

$$\alpha[a, b] + \beta[a, c] + \gamma[b, c] = 0.$$

If $\alpha = \beta = 0$ then $[b, c] = 0$ and we are done. If $\alpha = 0$ and $\beta \neq 0$, then $[\beta a + \gamma b, c] = 0$, and we replace a by $\beta a + \gamma b$. In the new generating set $[a, c] = 0$. Similarly, if $\alpha \neq 0$ and $\beta = 0$, then we replace a by $-\alpha a + \gamma c$. If $\alpha \neq 0$ and $\beta \neq 0$, we replace a by $(\beta/\alpha)a + (\gamma/\alpha)b$ and b by $b + (\beta/\alpha)c$. In the two latter cases, the new generating set satisfies $[a, b] = 0$. Thus after possibly reordering the generators a, b, c we may assume that $L_2 = \langle [a, b], [a, c] \rangle$, $[b, c] = 0$, and $L_4 = \langle [[a, b], [a, c]] \rangle$.

We claim that the elements a, b, c can be chosen so that, in addition to what was proved in the last paragraph, we also have $[a, b, a] = [a, b, b] = [a, c, a] = 0$. First we modify the chosen elements to make $[a, b, a]$ zero. If $[a, b, a] = 0$, then we have nothing to do, and if $[a, c, a] = 0$ then we interchange b and c . So assume that $[a, b, a] \neq 0$ and $[a, c, a] \neq 0$. Since L_3 is one-dimensional, we have that $\alpha[a, b, a] + \beta[a, c, a] = 0$ for some $\alpha, \beta \in \mathbb{F}_p$ with $\alpha\beta \neq 0$. Thus $[a, \alpha b + \beta c, a] = 0$ and we replace b by $\alpha b + \beta c$.

Since $[[a, b], [a, c]] = [a, b, a, c] - [a, b, c, a] = [a, b, c, a] \neq 0$ we have that $[a, b, c, a] \neq 0$. Thus a does not centralise L_3 . On the other hand, $[a, b, a, b] = [a, b, b, a] = 0$. If $[a, b, b] \neq 0$, then, as in this case $L_3 = \langle [a, b, b] \rangle$, we have that a centralises L_3 , which is not possible. Therefore $[a, b, b] = 0$ must hold. If $[a, c, a] \neq 0$ then, as $[a, b, c] = [a, c, b] \neq 0$ we have that

$$0 = \alpha[a, c, a] + \beta[a, c, b] = [a, c, \alpha a + \beta b]$$

with some $\alpha, \beta \in \mathbb{F}_p$ scalars such that $\alpha\beta \neq 0$. If we replace a by $\alpha a + \beta b$, then we obtain the additional relation $[a, c, a] = 0$. It is easy to see that the previously proved relations still hold for the new elements a, b, c .

Hence we proved that elements $a, b, c \in L_1$ can be chosen so that $L_2 = \langle [a, b], [a, c] \rangle$, $L_4 = \langle [[a, b], [a, c]] \rangle$, $[a, c] = [a, b, a] = [a, b, b] = [a, c, a] = 0$. Now

$$\begin{aligned} 0 &= [a, b, a, c] + [a, c, [a, b]] + [c, [a, b], a] \\ &= [a, b, a, c] + [a, c, a, b] - [a, c, b, a] - [a, c, b, a] = -2[a, c, b, a]. \end{aligned}$$

Thus $2[a, c, b, a] = 0$, and so

$$2[[a, b], [a, c]] = -2[a, b, c, a] = -2[a, c, b, a] = 0.$$

As $[[a, b], [a, c]] \neq 0$ we obtain that $p = 2$. This completes the proof of Theorem 1.2.

Next we study two immediate consequences of Theorem 1.2.

Lemma 5.1. *Let G be a finite p -group.*

- (i) *If, for $d \geq 1$, the quotient $G^{(d)}\gamma_{2^{d+1}}(G)/\gamma_{2^{d+1}}(G)$ is cyclic, then, for $e \geq 1$, we have $G^{(d+e)} \leq \gamma_{2^{d+e}+2^{e-1}}(G)$. Further, if $G^{(d+e+1)} \neq 1$, then*

$$\log_p |G^{(d+e)}/G^{(d+e+1)}| \geq 2^{d+e} + 2^{e-1} + 1.$$

- (ii) *If, for $d \geq 0$, the quotient $G^{(d)}\gamma_{2^{d+1}}(G)/\gamma_{2^{d+1}}(G)$ can be generated by two elements, then $G^{(d+1)}\gamma_{2^{d+1}+1}(G)/\gamma_{2^{d+1}+1}(G)$ is cyclic, and so, for $e \geq 2$, we have $G^{(d+e)} \leq \gamma_{2^{d+e}+2^{e-2}}(G)$. Further, if $G^{(d+e+1)} \neq 1$, then*

$$\log_p |G^{(d+e)}/G^{(d+e+1)}| \geq 2^{d+e} + 2^{e-2} + 1.$$

Proof. (i) If $G^{(d)}\gamma_{2^{d+1}}(G)/\gamma_{2^{d+1}}(G) \cong G^{(d)}/(G^{(d)} \cap \gamma_{2^{d+1}}(G))$ is cyclic, then, by Lemma 2.2,

$$G^{(d+1)} = [G^{(d)}, G^{(d)}] = [G^{(d)}, G^{(d)} \cap \gamma_{2^{d+1}}(G)] \leq [G^{(d)}, \gamma_{2^{d+1}}(G)] \leq \gamma_{2^{d+1}+1}(G).$$

Now easy induction implies our claim.

- (ii) Suppose that $G^{(d)}\gamma_{2^{d+1}}(G)/\gamma_{2^{d+1}}(G)$ is generated by $x\gamma_{2^{d+1}}(G)$ and $y\gamma_{2^{d+1}}(G)$. If $a, b \in G^{(d)}$ then there are some $\alpha_a, \beta_a, \alpha_b, \beta_b \in \mathbb{Z}$ and $u_a, u_b \in \gamma_{2^{d+1}}(G)$ such that $a = x^{\alpha_a}y^{\beta_a}u_a$ and $b = x^{\alpha_b}y^{\beta_b}u_b$

$$[a, b] = [x^{\alpha_a}y^{\beta_a}u_a, x^{\alpha_b}y^{\beta_b}u_b] \equiv [x, y]^{\alpha_a\beta_b - \beta_a\alpha_b} \pmod{\gamma_{2^{d+1}+1}(G)}.$$

Hence

$$\frac{G^{(d+1)}\gamma_{2^{d+1}+1}(G)}{\gamma_{2^{d+1}+1}(G)} \cong \frac{G^{(d+1)}}{G^{(d+1)} \cap \gamma_{2^{d+1}+1}(G)}$$

is cyclic, and so our assertion follows from (i). $\square$

Theorem 1.2 and the previous lemma imply the following corollary.

Corollary 5.2. *If p is an odd prime, G is a finite p -group, $d \geq 1$, and $G^{(d)}/G^{(d+1)}$ is a small derived quotient, then, for $e \geq 1$, we have $G^{(d+e)} \leq \gamma_{2^{d+e}+2^{e-1}}(G)$. Further, if $G^{(d+e+1)} \neq 1$, then*

$$\log_p |G^{(d+e)}/G^{(d+e+1)}| \geq 2^{d+e} + 2^{e-1} + 1.$$

Proof. Theorem 1.2 implies that $|G^{(d)}/[G^{(d)}, G]| = p$. Since $[G^{(d)}, G] \leq G^{(d)} \cap \gamma_{2^{d+1}}(G)$, we obtain that

$$\left| \frac{G^{(d)}\gamma_{2^{d+1}}(G)}{\gamma_{2^{d+1}}(G)} \right| = \left| \frac{G^{(d)}}{G^{(d)} \cap \gamma_{2^{d+1}}(G)} \right| \leq \left| \frac{G^{(d)}}{[G^{(d)}, G]} \right| = p.$$

Hence $G^{(d)}\gamma_{2^{d+1}}(G)/\gamma_{2^{d+1}}(G)$ is cyclic and the required result follows from Lemma 5.1. $\square$

6. TOWARDS THE PROOF OF THEOREM 1.1

The coefficient “3” of the linear term of the bound in Theorem 1.1 is a consequence of the following theorem.

Theorem 6.1. *Suppose that p is a prime different from 3, G is a finite p -group, $d \geq 0$, and that $G^{(d)}\gamma_{2^{d+1}}(G)/\gamma_{2^{d+1}}(G)$ is elementary abelian. If the generator number of $G^{(d+2)}\gamma_{2^{d+2}+1}(G)/\gamma_{2^{d+2}+1}(G)$ is at least 3, then either $|\gamma_{2^d}(G)/\gamma_{2^{d+1}}(G)| \geq p^4$ or $|\gamma_{2^{d+1}+2^d}(G)/\gamma_{2^{d+1}+2^{d+1}}(G)| \geq p^2$.*

The proof of this theorem will be presented at the end of this section.

Let G denote a finite p -group. Suppose that K is the subring corresponding to $G^{(d)}$ in the Lie ring associated with the lower central series of G . That is

$$K = \bigoplus_{i \geq 2^d} (G^{(d)} \cap \gamma_i(G))\gamma_{i+1}(G)/\gamma_{i+1}(G).$$

The first homogeneous component of K is $K_{2^d} = G^{(d)}\gamma_{2^{d+1}}(G)/\gamma_{2^{d+1}}(G)$. We let L denote the subring generated by K_{2^d} . For $i \geq 1$, we let $L_i = L \cap K_{i2^d}$. Since L is generated by homogeneous elements of the same degree, we obtain that the L_i form a grading on L .

Let me warn the reader that the Lie algebras K and L in the previous paragraph are different from those introduced in Section 4.

Lemma 6.2. *For $i \geq 1$, we have $L_i \cong \gamma_i(G^{(d)})\gamma_{i2^d+1}(G)/\gamma_{i2^d+1}(G)$. If the factor group $G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G)$ has exponent p , then L can be considered as a Lie algebra over $\mathbb{F}_p$. Further, if the generator number of $G^{(d+2)}\gamma_{2^{d+2}+1}(G)/\gamma_{2^{d+2}+1}(G)$ is at least 3, then the following hold:*

- (i) $\dim(L'' \cap L_4) \geq 3$;
- (ii) $\dim L_1 \geq 3$, $\dim L_2 \geq 3$, and $\dim L_3 \geq 1$.

Proof. Since L is degree-1 generated, we obtain that $L_i = [L_1, \dots, L_1]$ with L_1 occurring i times. Hence the definition of the product in L gives

$$\begin{aligned} L_i &= [\underbrace{L_1, \dots, L_1}_{i \text{ times}}] = [\underbrace{G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G), \dots, G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G)}_{i \text{ times}}] \\ &= [\underbrace{G^{(d)}, \dots, G^{(d)}}_{i \text{ times}}]\gamma_{i2^d+1}(G)/\gamma_{i2^d+1}(G) = \gamma_i(G^{(d)})\gamma_{i2^d+1}(G)/\gamma_{i2^d+1}(G). \end{aligned}$$

If $L_1 = G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G)$ has exponent p , then L_1 is a vector space over $\mathbb{F}_p$, and, as L is generated by L_1 , we obtain that L can be considered as a Lie algebra over $\mathbb{F}_p$.

Now assume that the generator number of $G^{(d+2)}\gamma_{2^{d+2}+1}(G)/\gamma_{2^{d+2}+1}(G)$ is at least 3.

- (i) As L is degree-1 generated,

$$\begin{aligned} L'' \cap L_4 &= [L_2, L_2] = [[L_1, L_1], [L_1, L_1]] \\ &= [[G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G), G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G)], \\ &\quad [G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G), G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G)]] \\ &= [[G^{(d)}, G^{(d)}], [G^{(d)}, G^{(d)}]]\gamma_{2^{d+2}+1}(G)/\gamma_{2^{d+2}+1}(G) = G^{(d+2)}\gamma_{2^{d+2}+1}(G)/\gamma_{2^{d+2}+1}(G). \end{aligned}$$

Recall that L_4 is a vector space and, by assumption, the generator number of the factor group $G^{(d+2)}\gamma_{2^{d+2}+1}(G)/\gamma_{2^{d+2}+1}(G)$ is at least 3. Therefore $\dim(L'' \cap L_4) \geq 3$.

(ii) Recall that L is generated by L_1 and that $\dim(L'' \cap L_4) = \dim[L_2, L_2] \geq 3$. If L_2 is 1-dimensional, then $[L_2, L_2] = 0$, and so this is impossible. If L_2 is 2-dimensional and $\{x, y\}$ is a basis for L_2 then $[L_2, L_2]$ is generated by $[x, y]$, and so $[L_2, L_2]$ is at most 1-dimensional. Hence L_2 is at least 3-dimensional. As L is degree-1 generated, we have $L_2 = [L_1, L_1]$, and so similar argument shows that L_1 is also at least 3-dimensional. Since $L_4 \neq 0$, we must have $L_3 \neq 0$, and so $\dim L_3 \geq 1$. $\square$

Theorem 6.1 is a simple consequence of the following result.

Lemma 6.3. *Let p be a prime different from 3. If $G^{(d)}\gamma_{2^d+1}(G)/\gamma_{2^d+1}(G)$ has exponent p and $G^{(d+2)}\gamma_{2^{d+2}+1}(G)/\gamma_{2^{d+2}+1}(G)$ is a 3-generator abelian group, then either $\dim L_1 \geq 4$ or $\dim L_3 \geq 2$.*

Proof. By Lemma 6.2, in this case, L is a Lie algebra over $\mathbb{F}_p$, $\dim L_1 \geq 3$, $\dim L_2 \geq 3$, $\dim L_3 \geq 1$, and $\dim(L_4 \cap L'') \geq 3$. Suppose that the assertion of this lemma is not valid, and so $\dim L_1 = 3$ and $\dim L_3 = 1$. Let $\{x, y, z\}$ is a basis of L_1 . The subspace $L_4 \cap L''$ is spanned by

$$[[x, y], [x, z]] = [x, y, x, z] - [x, y, z, x], \quad [[x, y], [y, z]] = [x, y, y, z] - [x, y, z, y],$$

and

$$[[x, z], [y, z]] = [x, z, y, z] - [x, z, z, y].$$

If $[x, y, x]$ and $[x, y, z]$ are both zero then $[[x, y], [x, z]] = 0$ and $\dim L_4 \cap L'' \leq 2$. Suppose therefore that at least one of them is non-zero.

Assume first that $[x, y, x] \neq 0$. Since $\dim L_3 = 1$ we must have that

$$L_3 = \langle [x, y, x] \rangle \quad \text{and} \quad L_4 = \langle [x, y, x, x], [x, y, x, y], [x, y, x, z] \rangle$$

and there cannot be any linear dependencies among the three generators of L_4 . There must be some α such that $[y, x, y] = \alpha[x, y, x]$. By the Jacobi identity

$$[y, x, [y, x]] + [x, [y, x], y] + [y, x, y, x] = 0,$$

that is $[x, y, x, y] + [y, x, y, x] = 0$. In other words $[x, y, x, y] = -\alpha[x, y, x, x]$ which means that $L_4 = \langle [x, y, x, x], [x, y, x, z] \rangle$ therefore $\dim L_4 \leq 2$, which is a contradiction.

Suppose now that $[x, y, z] \neq 0$. As before we must have that

$$L_3 = \langle [x, y, z] \rangle \quad \text{and} \quad L_4 = \langle [x, y, z, x], [x, y, z, y], [x, y, z, z] \rangle.$$

The third homogeneous component in the free Lie algebra with rank 3 has a basis formed by the following 8 elements:

$$[y, x, x], [y, x, y], [y, x, z], [z, x, x], [z, x, z], [z, y, x], [z, y, y], [z, y, z].$$

The relation $[z, x, y] = [y, x, z] + [z, y, x]$ also holds. Since the homogeneous component L_3 is spanned by $-[y, x, z] = [x, y, z]$, the rest of the basis elements can be expressed as

its scalar multiples:

$$\begin{aligned} [y, x, x] &= \alpha_1[x, y, z], & [y, x, y] &= \alpha_2[x, y, z], & [z, x, x] &= \alpha_3[x, y, z], \\ [z, x, z] &= \alpha_4[x, y, z], & [z, y, x] &= \alpha_5[x, y, z], & [z, y, y] &= \alpha_6[x, y, z], \\ [z, y, z] &= \alpha_7[x, y, z] \end{aligned}$$

where $\alpha_i \in \mathbb{F}_p$ for $1 \leq i \leq 7$. The Jacobi identity implies that $[y, x, x, y] - [y, x, y, x] = 0$. That is to say that $\alpha_1[x, y, z, y] - \alpha_2[x, y, z, x] = 0$. Therefore $\dim L_4 \leq 2$ unless $\alpha_1 = \alpha_2 = 0$. In the same way, $[z, x, x, z] - [z, x, z, x] = 0$ and $[z, y, y, z] - [z, y, z, y] = 0$ yield $\alpha_3 = \alpha_4 = 0$ and $\alpha_6 = \alpha_7 = 0$, respectively. It is also a consequence of the Jacobi identity that

$$[y, z, [y, x]] + [z, [y, x], y] + [y, x, y, z] = 0,$$

that is

$$[y, z, y, x] - [y, z, x, y] - [y, x, z, y] + [y, x, y, z] = 0.$$

In other words $[y, z, x, y] + [y, x, z, y] = 0$, hence

$$\alpha_5[x, y, z, y] + [x, y, z, y] = (\alpha_5 + 1)[x, y, z, y] = 0.$$

Similarly

$$[y, x, [x, z]] + [x, [x, z], y] + [x, z, y, x] = 0$$

and

$$[z, x, [z, y]] + [x, [z, y], z] + [z, y, z, x] = 0$$

yield

$$(\alpha_5 - 2)[x, y, z, x] = 0 \quad \text{and} \quad (2\alpha_5 - 1)[x, y, z, z] = 0,$$

respectively. Then it must follow that

$$\alpha_5 + 1 = 0, \quad \alpha_5 - 2 = 0 \quad \text{and} \quad 2\alpha_5 - 1 = 0$$

which implies that $p = 3$ and $\alpha_5 = 2$. □

The following example shows that the condition $p \neq 3$ is necessary in Lemma 6.3.

Example 6.4. If $p = 3$ then a computation with the LIENQ program (see Schneider [Sch97]) shows that in the Lie algebra

$$L = \langle x, y, z \mid [y, x, x], [y, x, y], [z, x, x], [z, x, z], [z, y, x] = 2[x, y, z], [z, y, y], [z, y, z] \rangle$$

we have that $\dim L_1 = 3$, $\dim L_2 = 3$, $\dim L_3 = 1$, $\dim L_4 \cap L'' = 3$.

We can now prove Theorem 6.1.

Proof of Theorem 6.1. By Lemmas 6.2 and 6.3, either

$$\left| \frac{\gamma_3(G^{(d)})\gamma_{2^{d+1}+2^{d+1}}(G)}{\gamma_{2^{d+1}+2^{d+1}}(G)} \right| \geq p^2 \quad \text{or} \quad \left| \frac{G^{(d)}\gamma_{2^{d+1}}(G)}{\gamma_{2^{d+1}}(G)} \right| \geq p^4.$$

Thus either $|\gamma_{2^{d+1}+2^d}(G)/\gamma_{2^{d+1}+2^{d+1}}(G)| \geq p^2$ or $|\gamma_{2^d}(G)/\gamma_{2^{d+1}}(G)| \geq p^4$. $\square$

Corollary 6.5. *Suppose that $p \neq 3$, G is a finite p -group, and $d \geq 2$ such that the generator number of $G^{(i)}\gamma_{2^{i+1}}(G)/\gamma_{2^{i+1}}(G)$ is at least 3 for all $i \leq d$. Then $\log_p |G/G^{(d)}| \geq 2^d + 3d - 2$ and $\log_p |G| \geq 2^d + 3d + 1$.*

Proof. Note that, for $i = 0, \dots, d$, the assumptions of the theorem imply that the order of $\gamma_{2^i}(G)/\gamma_{2^{i+1}}(G)$ is at least p^3 . By Theorem 6.1, for $i = 0, \dots, d-2$, if $\gamma_{2^i}(G)/\gamma_{2^{i+1}}(G)$ has exponent p , then either $|\gamma_{2^i}(G)/\gamma_{2^{i+1}}(G)| \geq p^4$ or $|\gamma_{2^{i+1}+2^i}(G)/\gamma_{2^{i+1}+2^{i+1}}(G)| \geq p^2$. If the exponent of $\gamma_{2^i}(G)/\gamma_{2^{i+1}}(G)$ is at least p^2 , then we also have $|\gamma_{2^i}(G)/\gamma_{2^{i+1}}(G)| \geq p^4$. Summing this up, the required result follows. $\square$

7. THE PROOF OF THEOREM 1.1

We conclude this paper with the proof of Theorem 1.1.

Suppose, in this section, that $p \geq 5$, G is a finite p -group, and d is an integer such that $G^{(d)} \neq 1$. By Mann's result [Man00], $\log_p |G| \geq 2^d + 2d - 2$, and so we may assume that $d \geq 5$.

Let k denote the smallest non-negative integer such that the generator number of $G^{(k)}\gamma_{2^{k+1}}(G)/\gamma_{2^{k+1}}(G)$ is less than 3. Thus the generator number of $G^{(i)}\gamma_{2^{i+1}}(G)/\gamma_{2^{i+1}}(G)$ is at least 3 for all $i \in \{0, \dots, k-1\}$, and, in particular, $G^{(k-1)}\gamma_{2^{k-1}+1}(G)/\gamma_{2^{k-1}+1}(G)$ has order at least p^3 . This also implies that $G^{(k-1)} \neq 1$, and so we may assume without loss of generality that $d \geq k-1$. If $k = 0$, then G is generated by two elements, and so, $\log_p |G^{(i)}/G^{(i+1)}| \geq 2^i + 2^{i-2} + 1$ for $i \in \{2, \dots, d-1\}$. As $G^{(d)} \neq 1$, we have $\log_p |G| \geq 2^d + 2^{d-2} + d - 1$. As $d \geq 5$, we obtain that the assertion of the theorem is true in this case. If $k = 1$, then, for $i = 3, \dots, d-1$, we have $\log_p |G^{(i)}/G^{(i+1)}| \geq 2^i + 2^{i-3} + 1$. Then $\log_p |G| \geq 2^d + 2^{d-3} + d$, and so the theorem is valid also in this case. If $k = 2$, then $\log_p |G^{(i)}/G^{(i+1)}| \geq 2^i + 2$ for $i = 0, 1$, $|G^{(2)}/G^{(3)}| \geq p^5$, $|G^{(3)}/G^{(4)}| \geq p^9$. However, by Corollary 5.2 we have that either $|G^{(2)}/G^{(3)}| \geq p^6$ or $|G^{(3)}/G^{(4)}| \geq p^{10}$. For $i \geq 4$ we have that $\log_p |G^{(i)}/G^{(i+1)}| \geq 2^i + 2^{i-4} + 1$. Thus $\log_p |G| \geq 2^d + 2^{d-4} + d + 2$. Therefore, in this case, we have nothing to prove. Hence we may assume without loss of generality that $k \geq 3$.

By Corollary 6.5,

$$\log_p |G/G^{(k-1)}| \geq 2^{k-1} + 3k - 5 \quad \text{and} \quad \log_p |G| \geq 2^{k-1} + 3k - 2.$$

Thus, if $d = k - 1$, then $\log_p |G| \geq 2^d + 3d + 1$, the result is valid.

Let $d \geq k$. It follows from Lemma 3.2(c) that if $\log_p |G^{(k-1)}/G^{(k)}| = 2^{k-1} + 1$, then $[G^{(k-1)}, G] = G^{(k-1)} \cap \gamma_{2^{k-1}+1}(G)$, and so

$$\frac{G^{(k-1)}\gamma_{2^{k-1}+1}(G)}{\gamma_{2^{k-1}+1}(G)} \cong \frac{G^{(k-1)}}{G^{(k-1)} \cap \gamma_{2^{k-1}+1}(G)} = \frac{G^{(k-1)}}{[G^{(k-1)}, G]}.$$

However, in this case, $|G^{(k-1)}/[G^{(k-1)}, G]| \leq p^2$ (see Lemma 3.2), and so we must have that $\log_p |G^{(k-1)}/G^{(k)}| \geq 2^{k-1} + 2$. Thus

$$\begin{aligned} \log_p |G/G^{(k)}| &= \log_p |G/G^{(k-1)}| + \log_p |G^{(k-1)}/G^{(k)}| \\ &\geq 2^{k-1} + 3k - 5 + 2^{k-1} + 2 = 2^k + 3k - 3. \end{aligned}$$

Thus, if $d = k$, then $|G^{(k)}| \geq p$, and we find

$$\log_p |G| = 2^k + 3k - 2 = 2^d + 3d - 2,$$

and the result is, again, valid.

Assume now that $d \geq k + 1$. As $\log_p |G^{(k)}/G^{(k+1)}| \geq 2^k + 1$, we have

$$\begin{aligned} \log_p |G/G^{(k+1)}| &= \log_p |G/G^{(k)}| + \log_p |G^{(k)}/G^{(k+1)}| \\ &\geq 2^k + 3k - 3 + 2^k + 1 = 2^{k+1} + 3k - 2. \end{aligned}$$

Therefore, if $d = k + 1$, then $\log_p |G| \geq 2^d + 3d - 4$, and so the theorem is valid also in this case.

Therefore we suppose that $d \geq k + 2$. In this case, $\log_p |G^{(k+1)}/G^{(k+2)}| \geq 2^{k+1} + 1$. However, as $k \geq 3$, by Corollary 5.2, either

$$\log_p |G^{(k)}/G^{(k+1)}| \geq 2^k + 2$$

or

$$\log_p |G^{(k+1)}/G^{(k+2)}| \geq 2^{k+1} + 2$$

Thus

$$\begin{aligned} \log_p |G/G^{(k+2)}| &= \log_p |G/G^{(k+1)}| + \log_p |G^{(k+1)}/G^{(k+2)}| \\ &\geq 2^{k+1} + 3k - 2 + 2^{k+1} + 2 = 2^{k+2} + 3k. \end{aligned}$$

Hence if $d = k + 2$, then $\log_p |G| \geq 2^{k+2} + 3k + 1 = 2^d + 3d - 5$. Therefore, in this case, the theorem is valid.

Assume now that $d \geq k + 3$. Corollary 5.2 yields $\log_p |G^{(k+2)}/G^{(k+3)}| \geq 2^{k+2} + 2$. Hence

$$\begin{aligned} \log_p |G/G^{(k+3)}| &= \log_p |G/G^{(k+2)}| + \log_p |G^{(k+2)}/G^{(k+3)}| \\ &\geq 2^{k+2} + 3k + 2^{k+2} + 2 = 2^{k+3} + 3k + 2. \end{aligned}$$

Hence if $d = k + 3$, then $\log_p |G| \geq 2^{k+3} + 3k + 3 = 2^d + 3d - 6$, and so the result holds.

If $d \geq k + 4$, then, by Lemma 5.2, $\log_p |G^{(i)}/G^{(i+1)}| \geq 2^i + 3$, for all $i \in \{k+3, \dots, d-1\}$. Thus in this case the theorem also holds.

ACKNOWLEDGMENT

Much of the research presented in this paper was carried out while I was a PhD student at The Australian National University. I am particularly grateful to my PhD supervisor, Mike Newman, for his continuous support.

REFERENCES

- [Bla58] N. Blackburn. On a special class of p -groups. *Acta Math.*, 100:45–92, 1958.
- [Bla87] Norman Blackburn. The derived group of a 2-group. *Math. Proc. Cambridge Philos. Soc.*, 101(2):193–196, 1987.
- [Bok71] L. A. Bokut'. On nilpotent Lie algebras. *Algebra i Logika*, 10:135–168, 1971.
- [Bur13] W. Burnside. On some properties of groups whose orders are powers of primes, *Proc. London Math. Soc.* (2), 11:225–243, 1913.
- [Bur14] W. Burnside. On some properties of groups whose orders are powers of primes (second paper), *Proc. London Math. Soc.* (2), 13:6–12, 1914.
- [ER00] Susan Evans-Riley. *On the Derived Length of Finite, Graded Lie Rings with Prime-Power Order, and Groups with Prime-Power Order*. PhD thesis, The University of Sydney, 2000.
- [ERNXX] Susan Evans-Riley and M. F. Newman. On the derived length of nilpotent graded Lie rings and groups with primer-power order. *In preparation*.
- [ERNS99] Susan Evans-Riley, M. F. Newman, and Csaba Schneider. On the soluble length of groups with prime-power order. *Bull. Austral. Math. Soc.*, 59(2):343–346, 1999.
- [GlaXX] S. P. Glasby. Solvable groups with a given solvable length, and minimal composition length. To appear in *J. Group Theory*.
- [GlaYY] S. P. Glasby. The shape of solvable groups with odd order. Submitted.
- [Hal34] P. Hall. A contribution to the theory of groups of prime-power order. *Proc. London Math. Soc.* (2), 36:29–95, 1934.

- [KLGP97] G. Klaas, C. R. Leedham-Green, and W. Plesken. *Linear pro- p -groups of finite width*. Springer-Verlag, Berlin, 1997.
- [LGM02] C. R. Leedham-Green and S. McKay. *The Structure of Groups of Prime Power Order*. Oxford University Press, 2002.
- [Man00] Avinoam Mann. The derived length of p -groups. *J. Algebra*, 224(2):263–267, 2000.
- [Sch97] Csaba Schneider. Computing nilpotent quotients in finitely presented Lie rings. *Discrete Math. Theor. Comput. Sci.*, 1(1):1–16 (electronic), 1997.
- [Sch00] Csaba Schneider. *Some results on the derived series of finite p -groups*. PhD thesis, The Australian National University, 2000.
- [Sch03] Csaba Schneider. Groups of prime-power order with a small second derived quotient. *J. Algebra*, 266(2):539–551, 2003.
- [SchXX] Csaba Schneider. Small derived quotients in finite p -groups. [arXiv.org/math.GR/0510223](https://arxiv.org/abs/math/0510223).
- [Sha95] A. Shalev. Finite p -groups. In *Finite and locally finite groups (Istanbul, 1994)*, pages 401–450. Kluwer Acad. Publ., Dordrecht, 1995.

INFORMATICS LABORATORY, COMPUTER AND AUTOMATION RESEARCH INSTITUTE, THE HUNGARIAN ACADEMY OF SCIENCES, 1518 BUDAPEST PF. 63, HUNGARY

E-mail address: `csaba.schneider@sztaki.hu`

WWW: `www.sztaki.hu/~schneider`